

JOURNAL OF APPROXIMATION THEORY **41**, 291–296 (1984)Convergence Rates of  $\alpha$ -Stable Difference Methods

E. GÖRLICH AND D. PONTZEN\*

*Lehrstuhl A für Mathematik, RWTH Aachen,  
Templergraben 55, 5100 Aachen, West Germany**Communicated by P. L. Butzer*

Received August 12, 1983

## 1. INTRODUCTION

The purpose of this note is to prove a quantitative Lax type theorem expressed in terms of  $\alpha$ -stability and  $\alpha$ -well-posedness. Here the term “quantitative” means that in the consistency hypothesis upon the difference method a rate of convergence is prescribed and that in the convergence result a rate is obtained corresponding to the degree of smoothness of the initial value. Theorems of this sort have been established in a general Banach space setting by Butzer, Dickmeis, Nessel, and others [2, 4, 5, 7]. For specific  $L_p$  spaces such theorems were proven by Peetre and Thomée [10].

Our objective here is to work with  $\alpha$ -well-posedness for  $0 \leq \alpha < 1$ , instead of the usual (strong) well-posedness, that is, to admit initial value problems

$$\frac{d}{dt}u = Au \quad (t > 0), \quad u(0) = f \in X, \quad (1.1)$$

on a Banach space  $X$ , with the closed linear operator  $A$  forming the infinitesimal generator of a semigroup  $\{E(t); t > 0\}$  of growth order  $\alpha$  in the sense of Da Prato [6]. Essentially, the latter property requires that the operator norm of  $E(t)$  satisfies

$$\|E(t)\| \leq Mt^{-\alpha} e^{\omega t} \quad (t > 0). \quad (1.2)$$

For  $\alpha > 0$  this is a weaker property than strong well-posedness, to which it reduces for  $\alpha = 0$ . Examples of initial value problems which are  $\alpha$ -well-posed for some  $\alpha > 0$  but not 0-well-posed are frequently met among systems of differential equations of the form (1.1) when the symbol of  $A$  is not a normal matrix (cf. [8]; there also is given a characterization of  $\alpha$ -well-posedness analogous to the Kreiss theorem).

\*This author was supported by the Deutsche Forschungsgemeinschaft Grant Go 261/4.

To  $\alpha$ -well-posedness there corresponds the property of  $\alpha$ -stability for a difference scheme  $\{E_k; 0 < k \leq k_0\}$  of bounded linear operators on  $X$ , depending continuously on  $k$ . This requires the  $n$ th iterate of  $E_k$  to satisfy

$$\|E_k^n\| \leq C(nk)^{-\alpha} e^{Knk} \quad (0 < k \leq k_0, n \in \mathbb{N}), \quad (1.3)$$

for certain constants  $C, K$ . Again see [8] for a characterization.

In Section 2 a quantitative Lax-type theorem on a general Banach space  $X$  will be given, and in Section 3 this will be made more precise by specializing  $X$  to the Lebesgue space  $L_p^N$ .

## 2. GENERAL THEOREM

A difference scheme is said to be consistent with (1.1) of order  $\varphi(k)$  on some linear subspace  $D$  of  $X$  if, for each  $T > 0$ , there is a constant  $C_0$  such that, for each  $g \in D$ ,

$$\| |E_k - E(k)| E(t) g \| \leq C_0 k \varphi(k) \|g\|_D \quad (0 < k \leq k_0, 0 \leq t \leq T). \quad (2.1)$$

Here  $E(0)$  denotes the identity operator,  $\|\cdot\|_D$  is some norm on  $D$ , and  $\varphi$  is an increasing function with  $\varphi(x) \rightarrow 0$  as  $x \rightarrow 0+$ .

We further use the (Peetre)  $K$ -functional, which is defined by

$$K(t, f; X, D) = \inf_{g \in D} \{ \|f - g\| + t \|g\|_D \}, \quad (2.2)$$

for  $f \in X, t > 0$ .

Our first result is

**THEOREM 1.** *Let the initial value problem (1.1) be  $\alpha$ -well-posed for some  $\alpha \in [0, 1)$ , and let  $\{E_k; 0 < k \leq k_0\}$  be a difference scheme which is consistent with (1.1) of order  $\varphi(k)$  on a subspace  $D$  of  $X$ . The following assertions are equivalent:*

- (a) *the difference scheme is  $\alpha$ -stable;*
- (b) *for each  $T > 0$  there is a constant  $C_1$  such that*

$$\|E_k^n f - E(nk) f\| \leq C_1 (nk)^{1-\alpha} K(nk\varphi(k), f; X, D)$$

*for  $0 < k \leq k_0, nk \leq T$ , and  $f \in X$ ;*

- (c) *for arbitrary  $T > 0$  there is a constant  $C_2$  such that*

$$\begin{aligned} \|E_k^n f - E(nk) f\| &\leq C_2 (nk)^{1-\alpha} \|f\|, & f \in X, \\ &\leq C_2 (nk)^{1-\alpha} \varphi(k) \|f\|_D, & f \in D, \end{aligned}$$

*for  $0 < k \leq k_0, nk \leq T$ .*

*Proof.* Assuming (a) to hold, let  $g \in D$ . Using (1.3), (2.1), it follows for  $0 < k \leq k_0$ ,  $n \geq 2$ , and  $nk \leq T$  that

$$\begin{aligned} \|E_k^n g - E(nk) g\| &\leq \sum_{j=0}^{n-1} \|E_k^{n-j-1}\| \| [E_k - E(k)] E(jk) g \| \\ &\leq C_0 k \varphi(k) \|g\|_D \left\{ 1 + Ck^{-\alpha} \sum_{j=0}^{n-2} (n-j-1)^{-\alpha} \right\} \\ &\leq C_0 k \varphi(k) \|g\|_D \left\{ 1 + Ck^{-\alpha} \frac{n^{1-\alpha}}{1-\alpha} \right\} \\ &\leq C'_0 (nk)^{1-\alpha} \varphi(k) \|g\|_D, \end{aligned}$$

for some constant  $C'_0$  which is independent of  $n, k$ , and  $g$ . The case  $n = 1$  being trivial, property (6) follows by observing that

$$\begin{aligned} \|E_k^n f - E(nk) f\| &\leq \inf_{g \in D} \{ \| (E_k^n - E(nk))(f - g) \| + \| E_k^n g - E(nk) g \| \} \\ &\leq \inf_{g \in D} \{ (Ce^{KT} + Me^{\omega T})(nk)^{-\alpha} \|f - g\| \\ &\quad + C'_0 (nk)^{1-\alpha} \varphi(k) \|g\|_D \} \\ &\leq C_1 (nk)^{-\alpha} K(nk \varphi(k), f; X, D). \end{aligned}$$

The implication (b)  $\Rightarrow$  (c) follows from the very definition (2.2). Let (c) be satisfied. Inserting (1.2) into  $\|E_k^n - E(nk)\| \leq C_2 (nk)^{-\alpha}$  we have

$$\|E_k^n\| \leq (C_2 + Me^{\omega T})(nk)^{-\alpha} \quad (0 < k \leq k_0, nk \leq T),$$

which implies (1.3), that is, (a), and the proof is complete. ■

It may be noted that in (b) and (c) the estimate on  $D$  can be replaced by

$$\|E_k^n f - E(nk) f\| \leq C_1 k \varphi(k) \{1 + (nk)^{-\alpha} (n-1)\} \|f\|_D,$$

which is more precise in case  $n = 1$  (cf. (2.1)).

### 3. LEBESGUE SPACE CASE

Let  $L_p^N$  denote the  $N$  fold Cartesian product of  $L_p(\mathbb{R}^d)$ ,  $1 \leq p \leq \infty$ , with norm

$$\|f\|_p = \sum_{j=1}^N \|f_j\|_p \quad (f = (f_1, \dots, f_N) \in L_p^N), \quad (3.1)$$

where, for an  $f_j \in L_p(\mathbb{R}^d)$ ,

$$\|f_j\|_p = \left( \int_{\mathbb{R}^d} |f_j(x)|^p dx \right)^{1/p}, \quad 1 \leq p < \infty,$$

$$= \operatorname{ess\,sup}_{x \in \mathbb{R}^d} |f_j(x)|, \quad p = \infty.$$

By  $W_{p,m}^\lambda$  we denote the  $N$  fold product of Sobolev spaces

$$W_{p,m}^\lambda = \sum_{j=1}^N W_{p,m_j}(\mathbb{R}^d),$$

where  $m = (m_1, \dots, m_N)$  and  $m_j \in \mathbb{N} = \{0, 1, 2, \dots\}$  for each  $j$ , the norms being defined by

$$\|f\|_{p,m} = \sum_{j=1}^N \|f_j\|_{p,m_j} \quad (f = (f_1, \dots, f_N) \in W_{p,m}^\lambda), \quad (3.2)$$

with

$$\|f_j\|_{p,m_j} = \sum_{|r| \leq m_j} \|D^r f_j\|_p \quad (f_j \in W_{p,m_j}(\mathbb{R}^d)).$$

Here  $D^r$  is the differential operator  $\partial^{(r)}/\partial x_1^{r_1} \cdots \partial x_d^{r_d}$  and  $|r| = r_1 + \cdots + r_d$ .

We further need the Besov spaces

$$B_p^{s,\lambda,N} = \prod_{j=1}^N B_p^{s_j,\lambda}(\mathbb{R}^d) \quad (s = (s_1, \dots, s_N), s_j > 0),$$

with norms

$$\|f\|_{B_p^{s,\lambda,N}} = \|f\|_p + \sum_{j=1}^N \sup_{t>0} t^{-s_j} \omega_{m_j}(t, f_j)_p \quad (0 < s_j < m_j),$$

$$\omega_{m_j}(r, f_j)_p = \sup_{0 < |h| \leq t} \left\| \sum_{l=0}^{m_j} \binom{m_j}{l} (-1)^{m_j-l} f_j(\cdot + lh) \right\|_p \quad (h \in \mathbb{R}^d). \quad (3.3)$$

These are intermediate between  $L_p^N$  and  $W_{p,m}^\lambda$ , that is,

$$B_p^{s,\lambda,N} = \prod_{j=1}^N (L_p(\mathbb{R}^d), W_{p,m_j}(\mathbb{R}^d))_{s_j, m_j, \lambda} \quad (0 < s_j < m_j),$$

cf. [3, Sect. 4.3.1] for details.

The following special case of Theorem 1 can be considered as an extension to  $\alpha > 0$  of a result of Peetre–Thomée [10], cf. also Brenner–Thomée–Wahlbin [1; Sect. 3.3].

**THEOREM 2.** *Let the initial value problem (1.1) be  $\alpha$ -well-posed on  $L_p^N$  for some  $\alpha \in [0, 1)$ , and let  $\{E_k; 0 < k \leq k_0\}$  be a difference scheme which is  $\alpha$ -stable on  $L_p^N$  and consistent with (1.1) of order  $\varphi(k)$  on  $W_{p,m}^N$ , for some multi-index  $m$ .*

*Given  $s = (s_1, \dots, s_N)$  with  $0 < s_j < m_j$  and  $T > 0$ , there is a constant  $C_3$  such that, for each  $f \in B_p^{s, \infty, N}$ ,  $0 < k \leq k_0$ ,  $nk \leq T$ , one has*

$$\|E_k^n f - E(nk) f\|_p \leq C_3 \max_{1 \leq j \leq N} \{(nk)^{(s_j/m_j) - \alpha} \varphi(k)^{s_j/m_j}\} \|f\|_{B_p^{s, \infty, N}}.$$

*Proof.* By Theorem 1 we have for each  $f \in L_p^N$ ,

$$\|E_k^n f - E(nk) f\|_p \leq C_1 (nk)^{-\alpha} K(nk \varphi(k), f; L_p^N, W_{p,m}^N).$$

In view of [3; (4.3.4)] (for  $p = \infty$ , see also [9]) and (3.1)–(3.3), there is a constant  $C_m$  which does not depend on  $t$ , such that

$$\begin{aligned} K(t, f; L_p^N, W_{p,m}^N) &= \inf_{g_j \in W_{p,m_j}(\mathbb{R}^d)} \left\{ \sum_{j=1}^N \|f_j - g_j\|_p + t \sum_{j=1}^N \|g_j\|_{p, m_j} \right\} \\ &= \sum_{j=1}^N K(t, f_j; L_p(\mathbb{R}^d), W_{p,m_j}(\mathbb{R}^d)) \\ &\leq C_m \sum_{j=1}^N [\min(1, t) \|f_j\|_p + \omega_{m_j}(t^{1/m_j}, f_j)_p] \\ &\leq C_m \sum_{j=1}^N [\min(1, t) + t^{s_j/m_j}] \|f\|_{B_p^{s, \infty, N}} \\ &\leq NC_m \max_{1 \leq j \leq N} [\min(1, t) + t^{s_j/m_j}] \|f\|_{B_p^{s, \infty, N}} \\ &\leq 2NC_m \max_{1 \leq j \leq N} t^{s_j/m_j} \|f\|_{B_p^{s, \infty, N}} \quad (t > 0). \end{aligned}$$

Setting  $t = nk\varphi(k)$  the assertion follows. ■

We finally note that Theorem 2 might also be written as an equivalence theorem if one adds the case  $s_j = 0$ , where  $\|f\|_{B_p^{s, \infty, N}}$  has to be replaced by  $\|f\|_p$ .

## REFERENCES

1. PH. BRENNER, V. THOMÉE, AND L. B. WAHLBIN, "Besov Spaces and Applications to Difference Methods for Initial Value Problems," Lecture Notes in Mathematics No. 434, Springer-Verlag, Berlin/New York, 1975.

2. P. L. BUTZER, The Banach–Steinhaus theorem with rates, and applications to various branches of analysis, in “General Inequalities 2” (E. F. Beckenbach, ed.), pp. 299–331, Birkhäuser, Basel, 1980.
3. P. L. BUTZER AND H. BERENS, “Semi-Groups of Operators and Approximation,” Springer-Verlag, Berlin/New York, 1967.
4. P. L. BUTZER, W. DICKMEIS, H. JANSEN, AND R. J. NESSEL, Alternative forms with orders of the Lax equivalence theorem in Banach spaces, *Computing* **17** (1977), 335–342.
5. P. L. BUTZER AND R. WEIS, On the Lax equivalence theorem equipped with orders, *J. Approx. Theory* **19** (1977), 239–252.
6. G. DA PRATO, Semigrupper di crescita  $n$ , *Ann. Scuola Norm. Sup. Pisa* **20** (1966), 753–782.
7. W. DICKMEIS AND R. J. NESSEL, Classical approximation processes in connection with Lax equivalence theorems with orders, *Acta Sci. Math. (Szeged)* **40** (1978), 33–48.
8. E. GÖRLICH AND D. PONTZEN, Alpha well-posedness, alpha stability, and the matrix theorems of H.-O. Kreiss, to appear.
9. H. JOHNEN, Inequalities connected with the moduli of smoothness, *Mat. Vesnik* **9** (1972), 289–303.
10. J. PEETRE AND V. THOMÉE, On the rate of convergence for discrete initial value problems, *Math. Scand.* **21** (1967), 159–176.